# LIMNOLOGY and OCEANOGRAPHY



Limnol. Oceanogr. 9999, 2021, 1–15 © 2021 The Authors. Limnology and Oceanography published by Wiley Periodicals LLC on behalf of Association for the Sciences of Limnology and Oceanography. doi: 10.1002/lno.12000

# Coral heat tolerance under variable temperatures: Effects of different variability regimes and past environmental history vs. current exposure

# Verena Schoepf<sup>1,2,3\*</sup> Hermione Sanderson,<sup>2,3,4</sup> Ellis Larcombe<sup>2,3,4,a</sup>

<sup>1</sup>Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

<sup>2</sup>Oceans Graduate School and UWA Oceans Institute, The University of Western Australia, Perth, Western Australia, Australia
 <sup>3</sup>ARC Centre of Excellence for Coral Reef Studies, The University of Western Australia, Perth, Western Australia, Australia
 <sup>4</sup>Department of Biosciences, Swansea University, Swansea, UK

# Abstract

Exposure to high-frequency temperature variability often but not always enhances coral heat tolerance, raising the question of whether this depends on the type of variability regime and past vs. current exposure. We collected corals from a macrotidal, highly fluctuating temperature environment and preconditioned them to either constant or variable daily temperatures for ~ 1.5 yr. Corals were then exposed to three new temperature variability regimes for ~1 month (constant control, symmetric variability, and tidal variability) to assess the effect of short-term environmental history, followed by a 12-d heat stress test. Measurements of visual coral health, photophysiology, photosynthesis, respiration, and calcification rates showed that preconditioning to constant vs. variable temperatures for 1.5 yr did not significantly impact coral physiology and heat tolerance. In contrast, environmental history experienced in the month prior to the heat stress test significantly influenced the physiological responses, with corals exposed to both types of variability having lower heat tolerance. Interestingly, corals in the tidal variability regime suffered greater health declines than in the symmetric variability regime although both treatments had the same cumulative heat exposure. Since heating rate and temperature amplitude were higher in the tidal variability regime (but time spent above the bleaching threshold was shorter), this suggests that short, extreme heat pulses may be more deleterious than longer but more moderate ones, though other factors likely also played a role. Overall, our findings demonstrate that daily temperature variability has significant potential to alter coral heat tolerance but only certain types of variability may enhance coral adaptive capacity.

Anthropogenic climate change is one of the biggest threats to coral reefs today. Ocean warming in combination with increasingly frequent and intense marine heatwaves (Frölicher et al. 2018; Oliver et al. 2018) has significantly contributed to the decline of coral reefs worldwide as the resulting mass bleaching events can lead to coral mortality on regional to global scales (Eakin et al. 2010; Hughes et al. 2018). In addition, rising atmospheric  $CO_2$  concentrations also result in ocean acidification, defined as a reduction in seawater pH, which lowers the calcification rates of many marine calcifiers (Kroeker et al. 2013), including reef-building corals (Chan and Connolly 2013; Kornder et al. 2018). Thus, there is an urgent need to identify the mechanisms that may help corals persist in a rapidly changing ocean.

Many tropical coral reefs occur in waters where environmental conditions are relatively stable throughout the year. Thus, corals have evolved narrow environmental tolerance limits and live, for example, close to their upper thermal thresholds (Coles and Jokiel 1977; Fitt et al. 2001). When temperatures exceed their local maximum monthly mean (MMM) temperatures by only  $1-2^{\circ}$ C, the vital symbiosis with dinoflagellate algae (family Symbiodiniaceae) breaks down in a

<sup>\*</sup>Correspondence: v.schoepf@uva.nl

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Additional Supporting Information may be found in the online version of this article.

<sup>&</sup>lt;sup>a</sup>Present address: Institute of Biomedical and Environmental Health, School of Health and Life Sciences, University of the West of Scotland, Paisley, UK

Author Contribution Statement: V.S. designed the study, conducted the fieldwork, and statistical analyses. H.S. and E.L. conducted the mesocosm experiment, seawater carbonate chemistry, and physiological analyses. V.S. led the writing of the article, with all authors contributing to the final manuscript and approving its final version.

process referred to as coral bleaching (Berkelmans and Willis 1999), which is defined as a significant loss of symbionts and/or photosynthetic pigments from the coral tissue. Since corals meet up to 90% or more of their energetic requirements from autotrophic carbon translocated by their symbionts (Muscatine et al. 1981), this results in significant resource limitation, compromised physiology (e.g., reduced photosynthesis and calcification rates) and possibly death (Brown 1997; Baker et al. 2008), unless they can rely on alternative carbon sources such as zooplankton or their own tissue reserves (Grottoli et al. 2006; Anthony et al. 2009).

In recent years, evidence has accumulated that some corals persist in reef environments characterized by strong environmental variability and that exposure to such fluctuations has the potential to enhance stress tolerance (Palumbi et al. 2014; Schoepf et al. 2015; Rivest et al. 2017). In particular, highfrequency temperature variability-as for example encountered on a daily basis in back-reef pools and intertidal/shallow reefshas been shown to promote heat tolerance and mitigate bleaching risk in a number of species and sites (Safaie et al. 2018). However, some studies have also demonstrated that temperature variability can have no or detrimental effects on coral heat tolerance (Putnam and Edmunds 2011; Schoepf et al. 2019; Klepac and Barshis 2020), yet we lack a mechanistic understanding of why that is. One possible explanation is that only certain types of temperature variability (i.e., certain combinations of magnitude, amplitude and frequency) enhance heat tolerance and that this also depends on the type of variability encountered in their native environment, that is, their environmental history. In addition, other environmental parameters such as light often covary with temperature and could therefore further modulate how temperature variability impacts coral heat tolerance. However, these mechanisms are currently poorly understood because only few studies to date have compared how different types of variability regimes differ in their potential to alter stress tolerance (Rivest et al. 2017).

To address these knowledge gaps, we used corals from a highly fluctuating, naturally extreme temperature environment to compare the effects of different temperature variability regimes on heat tolerance. The corals originated from the macrotidal Kimberley region in NW Australia where the world's largest tropical tides (up to 12 m) expose corals on shallow, intertidal reefs to short-term maximum temperatures of up to 38°C, daily temperature variability of up to 8°C, and frequent aerial exposure at low tide (Dandan et al. 2015; Schoepf et al. 2015). Long-term acclimatization and/or adaptation to these extreme conditions have resulted in enhanced heat tolerance, survival, and recovery capacity post bleaching (Schoepf et al. 2015; Le Nohaïc et al. 2017; Jung et al. 2021). However, these corals were nevertheless unable to increase their heat tolerance after 6 months of acclimation to 1°C warmer temperatures in the lab; furthermore, under these warming levels, daily temperature variability actually lowered their ability to cope with a heat stress event compared to corals exposed to constant

daily temperatures (Schoepf et al. 2019). This raises the question of how environmental history interacts with different variability regimes as drivers of coral heat tolerance.

We preconditioned corals to either constant or variable daily temperatures for ~ 1.5 yr to create corals with different environmental history. Corals were then exposed to three different temperature variability regimes for ~ 1 month, followed by a heat stress test, to determine if (1) long- or short-term thermal history has a greater influence on coral heat tolerance and (2) different types of temperature variability differ in their ability to alter heat tolerance. In addition, we explored the influence of light variability on coral responses to different temperature variability regimes.

### Materials and Methods

### Overview

We used corals from an intertidal, highly variable reef environment in NW Australia as a model species and took advantage of these corals being grown long term (~ 1.5 yr) in the lab under two different temperature variability regimes (constant daily temperature and 4°C daily temperature variability; Fig. 1) to assess how environmental history influences coral heat tolerance. This preconditioning phase (see details below) was part of another experiment designed to investigate whether daily temperature variability enhances the capacity of corals to acclimate to future ocean warming (Schoepf et al. 2019). Unexpectedly, Schoepf et al. (2019) found that daily temperature variability did not promote long-term acclimation to ocean warming but instead often had negative effects on coral heat tolerance and physiology. The authors speculated that this result could be related to the fact that the somewhat artificial, symmetric temperature variability treatments did not simulate the tidally induced strong coupling of temperature and light variability experienced by intertidal corals in situ (Dandan et al. 2015; Schoepf et al. 2015, 2020). We therefore conducted another experiment (see the Temperature variability experiment section; Fig. 1) to expose the corals with different environmental history to three different types of temperature variability (constant daily temperature, 4°C daily temperature variability, and "tidal" temperature variability), combined with a more realistic "tidal" light regime (Fig. 2b). The "tidal" variability treatment mimicked the rapid changes observed in both temperature and light during spring tidal cycles at the collection site. Subsequently, we assessed how acclimation to these treatments influenced coral heat tolerance in a 12-d heat stress test.

### Study species and collection site

Intertidal Acropora aspera (Dana, 1846) corals were collected from Shell Island (Shenton Bluff), Cygnet Bay, in the macrotidal Kimberley region of NW Australia. Collection permits were obtained from the Western Australia Department of Fisheries (exemption no. 2549, date of issue 03 March



**Fig. 1.** Experimental design. Key events and experimental phases are shown on the left, while the timeline with details on temperature and light conditions is shown on the right. Black and blue colors represent the different environmental history established during the preconditioning phase. Arrows indicate replicate fragments per parent colony (A–E).

2015). The Kimberley region features the world's largest tropical tides (up to 12 m, ~8 m at Shell Island), thereby exposing shallow coral reefs to extreme environmental conditions as well as strong daily fluctuations in many environmental parameters (Schoepf et al. 2015; Cornwall et al. 2018). Five visibly healthy coral colonies (labeled A, B, C, D, E) were collected in April 2016 from the intertidal environment at Shell Island (16°28'45.8"S, 123°2'41.3"E), where they regularly experience prolonged aerial exposure during spring low tides (up to hours), high light levels (up to ~ 2000  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>) and extreme daily temperature fluctuations of up to 7°C, with short-term maxima of up to 38°C (Dandan et al. 2015; Schoepf et al. 2015, 2020). In contrast to subtidal conspecifics, intertidal A. aspera have a naturally higher heat tolerance (Dandan et al. 2015; Schoepf et al. 2015) and survive/recover better from bleaching events (Le Nohaïc et al. 2017; Schoepf et al. 2020; Jung et al. 2021). Both intertidal and subtidal corals are dominated by symbionts from the genus Cladocopium (previously clade C; LaJeunesse et al. 2018) (Schoepf et al. 2015, 2020); however, symbiont community composition nevertheless differs between intertidal and subtidal corals and may be linked to their differential bleaching resilience (Jung et al. 2021). Monthly average temperatures at Shell Island range from ~  $25^{\circ}$ C to  $31^{\circ}$ C, and the bleaching threshold was experimentally established to be ~  $32^{\circ}$ C, approximately 1°C above local MMM sea surface temperatures (SST) (Schoepf et al. 2015). Corals were collected at least 10 m apart to avoid collecting clones.

Colonies were shipped live and submerged in water to the University of Western Australia and maintained in indoor, flow-through aquaria at the Watermans Bay seawater facility at ~ 29°C ( $\pm$  1°C) to facilitate recovery and acclimation to tank conditions. From mid-June until the end of July 2016, corals were kept at their normal seasonal temperatures, with temperatures adjusted twice a month to track the seasonal profile at their collection site (Table S1) as detailed in Schoepf et al. (2019). In July 2016, each colony was fragmented into two pieces of 5–10 cm which were glued onto prelabeled plastic tiles. Light was provided on a 12 : 12 h light : dark cycle, following a natural diurnal light cycle with gradual increases up to 560 µmol m<sup>-2</sup> s<sup>-1</sup> at noon (Ledzeal S150 Plus). Corals were fed twice a week with live brine shrimp. Further details on the feeding regime and mesocosm tank setup are given below.

### Preconditioning phase (01 August 2016–05 February 2018)

The preconditioning phase served to establish two sets of corals with different environmental history: either constant



**Fig. 2.** Overview of experimental treatments. (**a**) The three temperature variability treatments during the experiment (10 Apr–26 May 2018). For reference, the maximum monthly mean (MMM) temperature at the collection site is also shown. (**b**) Comparison of the light treatments during the preconditioning phase (01 Aug 2016–05 Feb 2018) vs. the experimental phase (incl. fragment preparation; 06 Feb–26 May 2018).

daily temperatures or 4°C daily temperature variability (Figs. S1, 2a), while still following the seasonal temperature profile of their Kimberley collection site (Table S1). Each parent colony had one fragment exposed to constant daily temperature and another one exposed to 4°C daily temperature variability. Importantly, variability treatments were designed such that daily average temperature was the same in both constant and variable treatments. From 01 August 2016 until 30 May 2017, the corals were part of the seasonal control treatments (referred to as "native controls") in the experiment described in Schoepf et al. (2019). Three of the five colonies (C-E) were exposed to elevated temperatures (~ 32.4°C) from 17 to 30 May 2017 as part of the stress test in Schoepf et al. (2019); however, immediately after the stress test, temperature was returned to ambient seasonal temperatures, thus allowing these colonies to recover for more than 8 months (01 June 2017-05 February 2018; see Supplement for more details). From 01 June 2017 until 05 February 2018, all colonies continued to be kept in the same tank setup and under constant vs. variable daily temperatures while still following the seasonal temperature profile (Table S1).

The mesocosm setup is described in detail in previous papers (Schoepf et al. 2018, 2019). Briefly, both constant and variable temperature treatments had two replicate tanks per

treatment. Coral fragments were maintained in 55-liter transparent plastic tanks where seawater was being replaced at a rate of ~  $0.5 \text{ Lmin}^{-1}$ . Water motion was provided using a submersible pump (Macro Aqua;  $3000 \text{ L h}^{-1}$ ). Temperature was maintained using titanium heaters (WeiPro; 500 or 1000 W) and controlled via ApexFusion software (Neptune Systems). The Apex temperature probes were calibrated 1-2 times a week using a high-precision thermometer (Fisher Scientific Traceable). Light was provided on a 12 : 12 h light : dark cycle (06:00-18:00 h) using 150 W LED lights (Ledzeal S150 Plus) with custom-designed LED arrangements and colors to ensure a light spectrum similar to shallow tropical reef environments. The lights were programed to follow a natural diurnal light cycle, with gradual increases up to 560  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup> at noon (measured using an Apogee MQ-200 cosine-corrected planar PAR-meter) (Fig. 2b). Relatively high maximum light levels were chosen because intertidal Kimberley corals regularly experience high light levels depending on tidal elevation, water clarity, and cloud cover (up to ~ 2000  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>); however, these extreme light levels are only experienced short-term during spring low tide, and therefore intermediate levels were used for daily exposure in this study. The incoming seawater was pumped from 12 m depth and filtered through three sand filters (~  $20 \,\mu m$  nominal size). Corals were fed twice a week with live brine shrimp. Approximately 2.5 g of brine shrimp eggs were hatched, and the stock solution with live nauplii was then equally divided among all tanks. HOBO v2 temperature loggers (Onset) were deployed in each tank and continuously recorded seawater temperature every 5 min.

### Temperature variability experiment (February–May 2018) Preparation of new coral fragments (06 February–09 April 2018)

After 1.5 yr in the mesocosms, the corals had grown significantly and could be fragmented to produce a greater number of new, small fragments. However, due to heater malfunctioning, not all initial parent colonies survived. At the beginning of February 2018, the remaining four large corals from the constant temperature treatment (colonies B, C, D, and E) and the remaining three large corals from the variable treatment (colonies A, C, and E) were split into six smaller fragments each of 3-4 cm in size. The new pieces were then glued to prelabeled plastic tiles and allowed to recover and grow out for 2 months until 09 April 2018. During this time, corals with a history of constant daily temperature continued to be kept under constant temperatures (six replicate tanks), whereas corals with a history of 4°C daily temperature variability continued to be kept under variable conditions (six replicate tanks). Temperatures from this point onward were kept at ~ 30°C since SST at the collection site stay at or above ~ 30°C from January through May (Table S1).

In addition, corals were now being acclimated to a new "tidal" light cycle that mimicked daily light regimes during spring tides at the collection site (Fig. 2b, *see* Supplement for details). Specifically, the "tidal" light cycle mimicked the strong increase in light level during the morning low tide, followed by a rapid drop in light levels when tide pools get flushed during the incoming tide and subsequent relatively low light levels throughout the afternoon due to high turbid-ity associated with strong tidal currents (Dandan et al. 2015; Schoepf et al. 2020). Corals continued to be fed twice a week with live brine shrimp.

# Acclimation to different temperature variability regimes (10 April–08 May 2018)

Corals with different environmental history were then acclimated for 4 weeks (10 April–08 May 2018) to three different temperature variability regimes: (1) constant daily temperature (~  $30.0^{\circ}$ C), (2) 4°C daily temperature variability (~  $30.0^{\circ}$ C ± 2°C), and (3) "tidal" daily temperature variability (Figs. 1, 2a). The "tidal" temperature variability treatment mimicked strong increases in temperature across the morning spring low tide, followed by a rapid return to ambient temperatures with the incoming tide (06:00–08:59 31.5°C, 09:00–10:59 34.0°C, 11:00–17:59 30.0°C, 18:00–21:59 29.0°C, 22:00–05:59 30.0°C; Fig. 2a). Importantly, treatments were designed such that daily average temperatures were almost

identical in all three treatments; in addition, they were designed such that daily cumulative heat stress exposure was almost identical in the variable and tidal treatment (Fig. 2a; Table S2). Each treatment had four replicate tanks. Two fragments per parent colony per environmental history were randomly distributed across the three temperature variability treatments, leading to all possible combinations of having experienced temperature variability in the past only (during the preconditioning phase), during the temperature variability experiment only, during both preconditioning and the experiment, or not at all (Fig. 1).

### Temperature ramp-up and heat stress test (09-26 May 2018)

After 4 weeks of acclimation to the three temperature variability treatments, two replicate tanks per treatment were assigned to a heat stress treatment, whereas the other two replicate tanks were assigned as controls and remained at ~ 30°C (Fig. 1). Temperature was gradually increased by 0.5°C every 2 d in the heat stress tanks over 6 d to reach the known bleaching threshold of ~ 32°C for these corals (Schoepf et al. 2015, 2018, 2019). After 5 d at ~ 32°C, temperatures in heat stress tanks were increased by another 0.5°C on 20 May 2018, and by another 0.5°C on 22 May 2018 because  $F_v/F_m$  values remained relatively stable (*see* below). Thus, corals in heat stress tanks were kept at ~ 33°C for the last 5 d of the heat stress test (22–26 May 2018), which resulted in a significant stress response (*see* below).

### Monitoring of treatment conditions

Water temperature was controlled using the Apex control system, as described above. HOBO Pro v2 temperature loggers (Onset) were deployed in each tank and continuously recorded seawater temperature every 15 min from the start of the acclimation phase until the end of the experiment (10 April–26 May 2018). To quantify heat stress, daily average temperatures were used to calculate degree heating days (DHDs) for the heat stress treatments (Maynard et al. 2008). Instead of long-term mean summer temperatures, a MMM value of 30.827°C was used to calculate DHD, as per previous work (Schoepf et al. 2020). Further details regarding monitoring of temperature, salinity, and seawater carbonate chemistry are provided in the Supplement.

### Physiological analyses Visual health status

Visual health (as well as partial mortality) was recorded using the CoralWatch<sup>®</sup> Coral Health Chart (Siebeck et al. 2006) at five time points during the experiment by the same observer: ~ 2 weeks prior to the start of the acclimation phase (27 March 2018), 2.5 weeks after the acclimation phase had started (26 April 2018), at the beginning of the temperature ramp-up phase (10 May 2018), after 9 d of heat stress (24 May 2018), and 3 d after the end of the heat stress test (29 May 2018). Corals were scored on the upper surface of the branches, with light intensity at the highest setting. Although

visual assessment of coral health is less sensitive than quantification of symbiont density and/or chlorophyll a (Chl a) content, we chose this method because it is nondestructive and thus allowed for repeated measurements. The brightness/saturation scale of the Coral Health Chart was developed and rigorously calibrated using analyses of symbiont density and Chl a content, such that a change of two units in brightness indicates a significant change in symbiont density and Chl a content and thus the bleaching state of corals (Siebeck et al. 2006).

### Photophysiology

Photophysiological performance was assessed via both the effective (light-adapted) and maximum (dark-adapted) quantum yield of photosystem II ( $\Delta F/F_m'$  and  $F_v/F_m$ , respectively), which is an indicator of photochemical efficiency and bleaching sensitivity (Warner et al. 1999). Measurements were taken before the acclimation phase started (09 April 2018), two times during the acclimation phase (16 and 30 April 2018) and on the last day of the temperature ramp-up (14 May 2018). During the heat stress test, photophysiology was measured almost daily (except on 15, 18, and 24 May 2018) but during the last 2 d (25 and 26 May), only  $F_v/F_m$  was measured since corals were used for respirometry during the day (see below).  $\Delta F/F_{m}$  was measured at 11:30 h when all temperature treatments were at the same temperature; during this time, light intensity was set to maximum levels  $(560 \,\mu\text{mol}\,\text{m}^{-2}\,\text{s}^{-1})$ .  $F_v/F_m$  was measured ~ 45 min after lights turned off to assess the photochemical efficiency in the darkadapted state. These data were then used to calculate the maximum excitation pressure over photosystem II (Qm) (Iglesias-Prieto et al. 2004), which is an indicator of symbiont performance at peak sunlight.  $Q_m$  was calculated as  $Q_m = 1 - (\Delta F/$  $F_{\rm m}')/(F_{\rm v}/F_{\rm m})$ , with values close to 1 indicating photoinhibition and values close to 0 indicating light-limitation of photosynthesis under maximum irradiance. All photophysiological measurements were made using a Diving-PAM underwater fluorometer (Walz) with the following settings: measuring light intensity = 3, saturation pulse intensity = 12, saturation pulse width = 0.8 s, gain = 4, and damping = 2. Measurements were made at a constant distance of 2 mm from the coral tissue, approximately 2-3 cm below the tip. Variable and constant temperature treatments were assessed at the same temperature (i.e., temperatures in variable tanks were set to the same temperature as in the constant tanks during this time; during the heat stress test, all heated tanks were at the same temperature and all control tanks were at the same but lower temperature).

### Metabolism

Net photosynthesis and respiration rates were measured using respirometry at the end of the heat stress test (25–29 May) following the methods of Schoepf et al. (2019). Briefly, corals were incubated in clear ~ 1 liter chambers under both light (560  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup>) and dark conditions for 1.3–2.4 h, with stir bars providing continuous water motion. Control

chambers only contained a stir bar and no coral. Heat-stressed corals were incubated at 33°C, whereas control corals were incubated at 30°C. Heat-stressed corals were incubated prior to the control corals to minimize the risk of mortality due to cumulative heat stress exposure. Incubations were timed such that the variable temperature treatments were naturally at the same temperature as the constant treatments. At the beginning and end of each incubation, oxygen (Orion Star A323 RDO/DO meter, Thermo Fisher Scientific), salinity (YSI 85), pH, and temperature (Schott handylab pH 12, SI Analytics) were measured. The volume of the incubation seawater within the chambers was measured by pouring the seawater into a graduated cylinder after all measurements were completed. Hourly oxygen data were converted from percentage of O2 saturation to  $\mu$ mol L<sup>-1</sup> seawater using the equations of Garcia and Gordon (1992) and normalized to surface area (see below). Given the 12 : 12 h light : dark regime, P/R ratios were calculated as 12 h of gross P (= net P + R) divided by 24 h of R.

### Calcification

Calcification rate was determined using the buoyant weighing technique (Jokiel et al. 1978) at the beginning and end of both the acclimation phase (including temperature ramp-up) and the heat stress test. The wet weight of the coral (minus the combined weight of the tile and epoxy) was converted to dry weight using the density of seawater based on measurements of salinity and temperature and the known density of aragonite (2.93 g cm<sup>-3</sup>). Calcification rates were normalized to surface area and calculated as daily rates (mg d<sup>-1</sup> cm<sup>-2</sup>). Surface area was calculated as per Schoepf et al. (2019) using the relationship between coral skeletal mass (dry weight) and surface area determined via computed tomographic scans of skeletons of various sizes from the same coral species from the same location.

### Statistical analyses

A three-way PERMANOVA was used to test whether preconditioning to either constant or variable temperatures for ~ 1.5 yr prior to the experiment impacted the physiological response to the three variability treatments and heat stress test. This was tested using the following five response variables measured during or at the end of the heat stress test: visual health score (after 9 d of heat stress),  $F_v/F_m$  (on the last day of the heat stress test), calcification, photosynthesis, and respiration rate. However, the effect of preconditioning was neither significant on its own, nor in any of the interactions with variability and heat stress (Table 2). Thus, data for the two preconditioning treatments were pooled for further analysis. Data were normalized prior to analysis due to different measurement scales. The PERMANOVA was performed using Euclidean distances, type III (partial) sums of squares, and 9999 permutations using the software Primer v7 with the PER-MANOVA+ add-on package.

		Constant		Vari	able	Tidal	
		Control	Heated	Control	Heated	Control	Heated
Acclimation phase	Mean	30.04		30.36		30.56	
	SEM	0.01		0.02		0.02	
	<i>n</i> meas	6578		2720		2720	
	<i>n</i> tanks	4		4		4	
Temperature ramp-up	Mean	30.36	30.87	30.56	31.10	30.50	31.17
	SEM	0.01	0.04	0.05	0.06	0.05	0.05
	n meas	576	830	576	576	576	576
	<i>n</i> tanks	2	2	2	2	2	2
Heat stress test	Mean	30.23	32.38	30.29	32.53	30.64	32.66
	SEM	0.01	0.03	0.03	0.04	0.03	0.04
	<i>n</i> meas	1420	1429	1429	1429	1429	1429
	<i>n</i> tanks	2	2	2	2	2	2
DHD		0	26.5	0	28.6	0.6	29.7

**Table 1.** Seawater temperature (°C) for all treatments during the experiment based on continuous logger measurements. Degree heating days (DHD) were calculated for the combined temperature ramp-up phase and heat stress test.

Univariate statistical analyses were run separately for the acclimation phase and the phase involving heat stress and were performed in RStudio, R version 4.0.3 (R Core Team 2020). For the acclimation phase, linear mixed effect (LME) models were computed using the *lme4* package (Bates et al. 2015) for health score,  $F_v/F_m$  and  $Q_m$ . Variability regime (three levels: constant, variable, and tidal) and time (2-3 time points) were included as fixed factors. Time was included as a fixed factor because we were interested in whether the response variables changed over time; therefore, we included fragment ID (42 levels) to account for repeated measurements of individual coral fragments. Parent colony (five levels) was also included as a crossed random factor to account for colony-specific variability in physiological responses. Since assumptions associated with LME models were not met for health score and transformations were not successful, we instead ran a generalized linear mixed effects model (GLMM, Laplace approximation) using the *lme4* package and the Gamma distribution family (link = identity). The same model structure was used as for the LME models.

For the temperature ramp-up and heat stress phase, the above LME models were extended to include heat (two levels: control and heated) as a fixed effect in addition to variability regime and time, while fragment ID and parent colony remained random factors. However, with the exception of calcification rate, some modifications were required to account for response variables violating the assumptions of normality and homogeneity of variance (health score,  $F_v/F_m$ ,  $Q_m$ ) or the model failing to converge (health,  $F_v/F_m$ ). For health score and  $F_v/F_m$ , GLMM models were computed (Gamma distribution family, link = identity) but time had to be converted to a random factor for the model to converge; thus, fragment ID was removed from the model and parent colony was the only

other random factor for these variables. We additionally computed endpoint effects to test how variability regime and heat affected health score and  $F_v/F_m$  on the last two time points of the experiment.  $Q_m$  was cube root transformed to meet LME assumptions. Finally, photosynthesis and respiration rates were only measured once at the end of the experiment; thus a LME model was computed with variability regime and heat as fixed factors and parent colony as a random factor. Further details are provided in the Supplement. *p*-values < 0.05 were considered significant.

### Results

### Treatment conditions during the experiment

Seawater temperature, cumulative heat stress and carbonate chemistry for the acclimation phase, temperature ramp-up, and heat stress test are summarized in Tables 1 and 2, respectively. Seawater pH throughout the experiment was generally lower than ambient seawater (*see* Supplement for more details), with average values of ~ 7.8–7.9 that corresponded to pCO<sub>2</sub> levels of 520–780  $\mu$ atm and an aragonite saturation state of 2.8–3.6 (Table S3).

### Multivariate analysis and the effect of preconditioning

Preconditioning to either constant or variable temperatures for ~ 1.5 yr prior to the experiment did not significantly influence the overall physiological response as this effect was neither significant on its own, nor in any of the interactions with variability regime (during the experiment) and heat stress (Table 2). Thus, data for the two preconditioning treatments were pooled for univariate analyses (*see* below). However, the PERMANOVA also revealed that both variability regime and heat stress, as well as their interaction, significantly impacted physiological

**Table 2.** Results from three-way PERMANOVA analysis testing for the effect of preconditioning on the physiological response to temperature variability regimes and heat stress. Precond., preconditioning to either constant or variable temperatures; Var., variability treatments during the acclimation phase and heat stress test (three levels: constant, variable, and tidal); Heat, heat stress (two levels: control and heated); df, degrees of freedom; SS, sum of squares; MS, mean sum of squares; *P*(perm), permutational *p*-value; Unique perms, how many unique values of the test statistic were obtained under permutation. Significant *p*-values are highlighted in bold.

Effect	df	SS	MS	Pseudo-F	P(perm)	Unique perms
Precond.	1	6.65	6.65	1.70	0.2234	9963
Var.	2	36.17	18.09	4.63	0.0033	9960
Heat	1	57.03	57.03	14.60	0.0001	9959
Precond. $\times$ var.	2	4.22	2.11	0.54	0.6969	9974
Precond. $\times$ heat	1	2.73	2.73	0.70	0.5463	9967
Var. $\times$ heat	2	22.56	11.28	2.89	0.0439	9965
Precond. $\times$ var. $\times$ heat	2	4.73	2.36	0.60	0.6683	9962
Residuals	30	117.20	3.91			
Total	41	257.75				



Days of experiment

**Fig. 3.** Coral health. (a) Visual health of *A. aspera* (mean  $\pm$  SEM) ~ 2 weeks prior to the start of the acclimation phase (preAcclim), 2.5 weeks after the acclimation phase had started (Acclim), at the start of the temperature ramp-up phase (Ramp), after 9 d of heat stress (Heat) and 3 d after the heat stress test (postHeat). (b) Number of colonies showing partial and/or full mortality during the last day of the temperature ramp-up (day 98), the heat stress test (days 99–110) and 3 d after the heat stress test (days 111–113). Numbers in the treatments without mortality are slightly offset from zero for visual purposes. Dotted vertical lines indicate the end of the heat stress test. Significant main effects are indicated in italics (*see* Tables S3, S4). *n.s.*, no significant main effects. Asterisks indicate significant differences between control and heated corals in each variability (= var) regime, whereas  $\dagger$  indicates a significant difference between heated variable and heated tidal corals.

responses (Table 2). While heat stress did not significantly affect the physiology of corals maintained at constant temperatures during the experiment (t = 1.26, df = 10, p = 0.20), this was the case for corals in both the variable (t = 2.07, df = 10, p = 0.02) and tidal regime (t = 3.68, df = 10, p = <0.01).

### Coral health and physiology during the acclimation phase

Coral health score was high (~ 5) and stable during the acclimation phase and was also not affected by variability

regime (Fig. 3a; Table S4). The photochemical efficiency ( $F_v/F_m$ ) significantly increased over time (+ 8%) but this effect also depended on variability regime (Fig. 4a; Table S4). Prior to the acclimation phase, there were no significant differences between treatments but during the second half of the acclimation phase, both constant and tidal corals had higher  $F_v/F_m$  than the variable corals (+4% and +6%, respectively). The excitation pressure over PSII ( $Q_m$ ) was overall low (< 0.1) during the acclimation phase and, in contrast to  $F_v/F_m$ , stable over time,



**Fig. 4.** Photophysiology (mean  $\pm$  SEM). (a) Photochemical efficiency ( $F_v/F_m$ ) and (c) maximum excitation pressure over photosystem II ( $Q_m$ ) of *A. aspera* at various time points of the experiment. (b, d) Inserts show differences between control and heated corals during the temperature ramp-up phase and heat stress test. Significant main effects are indicated in italics (*see* Tables S3, S4). Asterisks indicate significant differences between control and heated corals in each variability (= var) regime, whereas  $\dagger$  indicates a significant difference between heated variable and heated tidal corals. Symbols indicate which variability treatments differed significantly from each other (independent of heat). Post hoc tests for the heat phase were only conducted for the last two measurement time points.

but corals maintained at constant temperatures had significantly higher (+60%)  $Q_{\rm m}$  than corals in the variable treatment (Fig. 4b; Table S4).

### Coral health and physiology during the temperature rampup and heat stress phase

Coral health score was significantly affected by both variability regime and an interactive effect of variability and heat as heated corals in the variable and tidal regime increasingly lost their color (i.e., bleached) during the heat stress test (Fig. 3a; Table S5). After 9 d of heat stress (day 108), only heated tidal corals had significantly lower health scores (-30%) than their respective controls, whereas 3 d after the heat stress test had ended (day 113), this was also the case for heated variable corals (-28% and -43% for variable and tidal corals, respectively). In contrast, heated corals maintained under constant temperatures did not suffer from a significant loss of color at any time point. Coral mortality mirrored these patterns (Fig. 3b), with heated tidal corals dying earlier (from day 103) and at higher rates (n = 5 fragments) than heated

variable corals (mortality from day 110, n = 3 fragments). No mortality occurred in any of the other treatments.

As for health score, significant effects of variability regime were observed for  $F_v/F_m$  but they depended on heat (Fig. 4a; Table S5). As heat stress progressed, heated corals in the variable and tidal regime increasingly experienced declines in  $F_v$ /  $F_{\rm m}$  whereas this was not the case for constant heated corals. However, temporal trends differed slightly between these groups, with heated variable corals showing the earliest reductions in  $F_v/F_m$ . After the first week of the heat stress test, however, they were surpassed by heated tidal corals and after 10 d of heat stress (day 109), heated tidal corals not only had significantly lower  $F_v/F_m$  than their controls (-19%) but also significantly lower  $F_v/F_m$  than the heated variable corals (-8%). Nevertheless, 1 d later on the last day of heat stress (day 110), this difference was no longer significant, with heated variable and tidal corals now having 17% and 21% lower  $F_v/F_m$  than their respective control group.

 $Q_{\rm m}$  showed a significant effect of variability regime and time; however, all two-way interaction terms including those

involving heat were also significant (Fig. 4b; Table S5). No significant differences between any treatments were observed during the first 5 d of the heat stress test but on day 6, heated corals in the tidal variability regime had significantly higher  $Q_{\rm m}$  (+84%) than heated variable corals (though  $Q_{\rm m}$  was not significantly higher than in the tidal controls).  $Q_{\rm m}$  of heated tidal corals then continuously increased throughout the heat stress test (up to ~ 0.22), being 86% higher than in the tidal controls and 71% higher than in the heated variable corals after 9 d of heat stress (endpoint for  $Q_{\rm m}$ ). In contrast, neither heated variable nor heated constant corals ever showed significantly higher or lower  $Q_{\rm m}$  than their respective controls at any point during the heat stress test.

Calcification rates were significantly influenced by variability regime, heat, and time; however, both the effect of variability regime and heat depended on the respective time point (Fig. 5; Table S5). During the combined acclimation and temperature ramp-up phase, neither heat nor variability regime significantly impacted calcification rates. However, during the heat stress phase, heated corals had significantly lower calcification rates (-83%) than the controls. Furthermore, corals maintained under the tidal variability regime had significantly lower calcification rates than corals maintained under either constant temperatures or in the variable treatment (-89% and -84%, respectively). This was largely due to heated tidal corals exhibiting negative calcification rates, that is, dissolution—the only treatment to do so.

The effects of variability regime and heat differed for photosynthesis and respiration rates as well as P/R ratios, and heat tended to have a greater effect than variability regime (Fig. 6; Table S6). Heated corals generally had significantly lower photosynthesis (-57%) but not respiration rates, resulting in 59% lower P/R ratios. Variability regime did not significantly affect



**Fig. 5.** Calcification (mean  $\pm$  SEM). Calcification rates of *A. aspera* during the combined acclimation and temperature ramp-up phase as well as during the heat stress test. Significant main effects are indicated in italics (*see* Table S5). Asterisks indicate significant differences between control and heated corals in each variability (= *var*) regime.

either photosynthesis or respiration rates (although a weak interactive effect of variability and heat was detected for *R*, p = 0.044); however, corals in the tidal variability regime had significantly lower *P*/*R* ratios (-25%) than corals maintained under constant daily temperatures and this effect was independent of heat (Fig. 6b). Overall, heated tidal corals had the lowest *P*/*R* ratios of all treatments, barely exceeding the threshold where corals transition from net autotrophy (*P*/*R* > 1) to net heterotrophy (*P*/*R* < 1).

### Discussion

This study provides new insights into the effects of environmental history on coral stress tolerance and is one of the first to compare how different temperature variability regimes differ in their ability to alter heat resistance over two temporal scales in an intertidal coral population. Specifically, we found that daily temperature variability only altered heat tolerance when experienced in the recent past (~ 1 month vs. ~ 1.5 yr) and that it generally lowered heat tolerance compared to exposure to constant daily temperatures.

### Effects of thermal history across two temporal scales

Although there is an increasing body of work demonstrating that a history of environmental variability has significant potential to alter coral resistance to climate change stressors (Rivest et al. 2017; Safaie et al. 2018; Kapsenberg and Cyronak 2019), few studies to date have specifically investigated the mechanisms and time scales over which environmental variability shapes stress tolerance. Many studies take advantage of natural gradients in environmental variability to compare the stress tolerance of coral populations along this gradient (Oliver and Palumbi 2011; Schoepf et al. 2015; Camp et al. 2016) but this approach, while powerful in detecting overall effects of environmental history, does not allow for distinguishing the underlying mechanisms. For example, environmental history may influence stress tolerance via longterm acclimatization and/or adaptation, environmental conditions experienced in the recent past (i.e., short-term acclimatization), carry-over effects across life stages or transgenerational acclimatization (Rivest et al. 2017), each mechanism acting over different temporal scales. In contrast, studies exposing corals to novel temperature conditions in the lab (Mayfield et al. 2012; Barshis et al. 2018; Schoepf et al. 2019) as well as reciprocal transplant experiments in combination with subsequent heat challenges (Palumbi et al. 2014; Klepac and Barshis 2020) enable us to better distinguish between shortand long-term acclimatization or adaptation or recent vs. native environment as dominant drivers of stress tolerance.

In the present study, preconditioning intertidal corals to either constant temperatures or  $4^{\circ}$ C daily temperature variability for ~ 1.5 yr had no effect on coral tolerance to a subsequent heat stress test. This finding is largely consistent with the lack of such an effect for this study species when



**Fig. 6.** Photosynthesis and respiration (mean  $\pm$  SEM). (**a**) Net photosynthesis (*P*) and dark respiration (*R*) rates as well as (**b**) *P*/*R* ratios of *A. aspera* at the end of the heat stress test. Dotted line in (**b**) highlights the transition from net autotrophy (*P*/*R* > 1) to net heterotrophy (*P*/*R* < 1). Significant main effects are indicated in italics (*see* Table S6). Asterisks indicate significant differences between control and heated corals in each variability (= *var*) regime, whereas  $\dagger$  indicates a significant difference between heated variable and heated tidal corals. Letters indicate which variability treatments differed significantly from each other (independent of heat).

preconditioning occurred over only ~ 9 months (Schoepf et al. 2019) (see Table S7 for how this study compares to previous work investigating the effects of temperature variability in corals from the macrotidal Kimberley region). Similarly, preconditioning Porites lobata corals from locations with differing levels of thermal variability to low and high thermal variation in common garden aquaria had minimal effects on their physiology and heat tolerance, although preconditioning only occurred for ~ 1 month (Barshis et al. 2018). Given that corals in the present study originated from an intertidal, highly variable temperature environment (up to 7°C daily temperature variability), it was expected that corals exposed to constant temperatures would have similar heat tolerance as the corals maintained under variable temperatures if longterm acclimatization and/or adaptation to high temperature variability at their native reef is the main mechanism shaping heat tolerance (Dixon et al. 2015; Kenkel et al. 2015; Barshis et al. 2018). In contrast, we expected them to have lower heat tolerance if it is primarily recent environmental conditions that shape future responses to heat stress (i.e., short-term acclimatization). For example, it has been shown that transplantation from highly variable to moderately variable temperature environments can lead to reduced heat tolerance in Acropora hyacinthus over 12-27 months (Palumbi et al. 2014). The lack of an observed effect of preconditioning suggests that intertidal A. aspera corals from the Kimberley region are long-term acclimatized or locally adapted to highly variable temperatures, which has led to enhanced bleaching resilience compared to conspecifics from the subtidal where daily temperature variability is much more moderate (Schoepf et al. 2020; Jung et al. 2021) (Table S7).

In contrast to the lack of an observed effect of long-term preconditioning, we found that thermal history experienced in the month prior to a heat stress event significantly altered bleaching sensitivity. This finding is in general agreement with several other laboratory studies investigating the effects of thermal history over similar time scales (i.e., days to ~1 month). For example, it has been shown that different heating rates (Middlebrook et al. 2010), thermal priming via short heat pulses (Bellantuono et al. 2012; Middlebrook et al. 2012; Ainsworth et al. 2016), or exposure to variable temperatures (Putnam and Edmunds 2011; Bay and Palumbi 2015) significantly influenced physiological responses and heat tolerance to subsequent heat stress. However, it is important to note that the majority of these studies observed that prior exposure to stressful temperatures or temperature variability improved heat tolerance (but see Putnam and Edmunds 2011; Schoepf et al. 2019; Klepac and Barshis 2020), whereas we observed the opposite (see below) and some studies also found no effect (Middlebrook et al. 2012; Barshis et al. 2018). Nevertheless, the results from this study highlight that recent thermal history can play an important role in shaping heat tolerance even when corals are long-term acclimatized or adapted to variable temperatures, such as in these intertidal corals from a highly fluctuating, macrotidal reef environment.

# Daily temperature variability lowered coral heat tolerance and survival

Although a recent meta-analysis has shown that high-frequency temperature variability increases coral heat tolerance and lowers bleaching risk (Safaie et al. 2018), we found the opposite pattern as daily temperature variability of  $4-5^{\circ}$ C

significantly enhanced sensitivity to heat stress and mortality risk in both the variable and tidal regime. This is particularly highlighted by the fact that exposure to heat stress equivalent to ~ 27 DHDs in corals maintained at constant daily temperatures elicited only minor negative responses as they did not suffer significant color loss or mortality and maintained  $F_{\rm v}/F_{\rm m}$ and calcification rates (but they had 45% lower P/R ratios). While we observed a similar negative effect of temperature variability on heat tolerance of intertidal A. aspera in a previous lab experiment (Schoepf et al. 2019), it contradicts our findings of enhanced bleaching resilience for this coral population in the field (Schoepf et al. 2015; Le Nohaïc et al. 2017; Jung et al. 2021) (Table S7). So how can we reconcile these seemingly contradictory findings, and why does highfrequency temperature variability not always promote heat tolerance, as also observed in some other studies (Putnam and Edmunds 2011; Klepac and Barshis 2020)?

We propose that a specific exposure threshold exists that determines whether high-frequency temperature variability is beneficial and enhances heat tolerance or results in chronic stress that exacerbates cellular damage and reduces overall resilience. As a result, only certain variability exposure regimes would be beneficial, potentially explaining the range of responses observed to date. This proposed variability threshold likely depends on several factors that are cumulative in nature and, just like the existence of such a threshold itself, are currently poorly understood. Important factors likely include amplitude and duration, as it is well-established for coral bleaching thresholds and the degree heating week (DHW) measure of cumulative heat stress (Liu et al. 2014), frequency (Safaie et al. 2018), and heating rate (Middlebrook et al. 2010) as well as the relationship between local variability regime and the characteristics of a given stress trajectory (Rivest et al. 2017).

The existence of such a variability exposure threshold is supported by several lines of evidence. First, several studies that observed positive effects of daily temperature variability on coral heat tolerance only exposed corals to short periods of heat stress (8 h to 5 d), thus mimicking tidal heating and short heatwave events rather than the prolonged heat stress that typically causes bleaching events (Bay and Palumbi 2015; Barshis et al. 2018). If the heat stress test in this study had only lasted for 5 d, negative effects of daily temperature variability would not have been evident (based on measurements of  $F_v/F_m$  and  $Q_m$ , Fig. 4). Second, Klepac and Barshis (2020) also found reduced thermal tolerance in massive corals from a highly variable back-reef pool which contradicted previous work demonstrating enhanced heat tolerance in these corals (Barshis et al. 2010, 2018). This unexpected finding was attributed to a thermal anomaly (5-8 DHW) during the 12 months reciprocal transplant experiment that preceded the heat stress assay and may have exceeded the upper thermal limits of native P. lobata corals.

Third, this study showed that two different variability regimes with almost identical cumulative heat exposure on

both a daily level (Tables 1, S2) and during the heat stress test (Table 1) differentially impacted coral heat tolerance. While both variability regimes increased sensitivity to heat stress, negative impacts were much more pronounced in the tidal variability regime for all responses measured. For example, tidal heated corals were the only treatment group to experience skeletal dissolution (Fig. 5) and P/R ratios indicating a transition from net autotrophy to net heterotrophy  $(P/R \sim 1)$ ; Fig. 6). It is possible that the more pronounced negative impacts on coral health were, at least in part, due to the fact that both daily average temperatures during the acclimation phase (30.56°C vs. 30.36°C) and cumulative heat exposure during the heat stress test (29.7 vs. 28.6 DHDs) were minimally higher in this treatment compared to the heated variable corals. However, given the small temperature differences, it seems more likely that other factors also played a role. For example, heated tidal corals spent less time daily at temperatures above the local bleaching threshold of ~ 31°C than the heated variable corals (5 vs. 9 h) but maximum daily temperatures were up to 2°C higher (34°C vs. 32°C), minimum daily temperatures were also higher (29°C vs. 28°C), they spent less time at the minimum daily temperatures (4 h at 29°C vs. 9 h at 28–29°C) and hourly heating rates were also higher (Fig. 2a; Table S2). In addition, maximum daily light levels were intentionally aligned with maximum daily temperatures in the tidal regime, whereas this was not the case in the variable regime, likely increasing overall stress in the heated tidal corals.

Finally, it is important to note that while the tidal regime mimicked the local variability regime at the collection site better than the variable regime (see Methods), the conditions simulated in the tidal regime are representative of spring rather than neap tides and thus the corals do not experience this kind of variability in their native environments for weeks on end. The acclimation phase demonstrated that intertidal corals can apparently cope with such extreme "spring-tide" temperature variability on a daily basis for prolonged periods of time, but suffer severe health declines and reduced heat tolerance when heat stress is super-imposed. The fact that we observed such contradictory findings of how strong daily temperature variability alters coral heat tolerance in this coral population (Table S7) further suggests the existence of a variability exposure threshold that is likely strongly dependent on the relationship between local variability regime and the characteristics of a given stress trajectory (Rivest et al. 2017).

Taken together, these lines of evidence suggest that (1) only certain variability exposures enhance coral stress tolerance depending on whether a time-integrated, cumulative exposure threshold is exceeded; (2) even when daily average temperatures and DHDs are very similar, stressor amplitude, time spent at both maximum and minimum daily temperatures (Mayfield et al. 2013), heating rate (Middlebrook et al. 2010), and covarying stressors play a key role in determining whether high-frequency temperature variability increases or lowers coral heat tolerance; and (3) when prolonged heat stress is

superimposed on an already highly variable temperature environment, protective mechanisms that enhance heat tolerance may quickly be overwhelmed—in some cases more easily or rapidly than in corals originating from or acclimated to more stable thermal environments (this study, Schoepf et al. 2019; Klepac and Barshis 2020).

Our findings provide new insights into how environmental history and different temperature variability regimes impact coral heat tolerance but it is important to note that seawater pH during the experiment was ~ 7.8–7.9 and thus lower than current ambient seawater pH (see Supplement for more details). We consider it unlikely that this significantly influenced coral heat tolerance because several independent studies have shown that bleaching sensitivity is not affected by seawater pH in six different coral species, including two Acropora species (e.g. Wall et al. 2013; Noonan and Fabricius 2015; Horvath et al. 2016). Importantly, one of these studies also mimicked daily temperature and pH variability in their treatments (van der Zande et al. 2020) and all tanks in our experiment received the same seawater, so that pH was similar across treatments (Table S3). Nonetheless, low seawater pH can impact several aspects of coral physiology, especially calcification (Chan and Connolly 2013; Kornder et al. 2018); therefore, when assessing absolute values of our response variables rather than comparing relative differences across treatments, this needs to be taken into account.

### Conclusion

In summary, our study shows that a history of thermal variability had a greater effect on coral heat tolerance when experienced in the recent rather than distant past (~ 1 month vs.  $\sim 1.5$  yr), at least in a population originating from a highly fluctuating, intertidal reef habitat. Further research is needed to determine whether corals from thermally less variable habitats, such as subtidal corals, show similar responses. In addition, our findings add to an increasing body of work demonstrating that strong daily temperature variability (4-5°C) can significantly lower coral heat tolerance when prolonged heat stress (12-18 d) is super-imposed on an already highly variable temperature regime (Putnam and Edmunds 2011; Schoepf et al. 2019; Klepac and Barshis 2020). While reef habitats with highly variable temperatures may select more effectively for heat-resistant genotypes than those with moderate variability (Palumbi et al. 2014; Kenkel et al. 2015; Schoepf et al. 2015), it is precisely the most variable habitats that may be at greatest risk during severe heat stress events because the combination of elevated background temperatures and natural variability may drive daily temperature maxima beyond sublethal thresholds (Klepac and Barshis 2020). Future research should therefore focus on identifying potential variability exposure thresholds as well as optimal variability characteristics such as amplitude, duration, and frequency to increase our understanding of when natural

variability will mitigate or amplify bleaching and mortality risk.

### Data availability statement

Data will be made available after publication in the online open access data repository Figshare.

### References

- Ainsworth, T. D., S. F. Heron, J. C. Ortiz, P. J. Mumby, A. Grech, D. Ogawa, C. M. Eakin, and W. Leggat. 2016. Climate change disables coral bleaching protection on the Great Barrier Reef. Science (80-) **352**: 338–342. doi:10. 1126/science.aac7125
- Anthony, K. R. N., M. O. Hoogenboom, J. A. Maynard, A. G. Grottoli, and R. Middlebrook. 2009. Energetics approach to predicting mortality risk from environmental stress: A case study of coral bleaching. Funct. Ecol. 23: 539–550. doi:10. 1111/j.1365-2435.2008.01531.x
- Baker, A. C., P. W. Glynn, and B. Riegl. 2008. Climate change and coral reef bleaching: An ecological assessment of longterm impacts, recovery trends and future outlook. Estuar. Coast. Shelf Sci. 80: 435–471. doi:10.1016/j.ecss.2008. 09.003
- Barshis, D. J., C. Birkeland, R. J. Toonen, R. D. Gates, and J. H. Stillman. 2018. High-frequency temperature variability mirrors fixed differences in thermal limits of the massive coral *Porites lobata*. J. Exp. Biol. **221**: jeb188581. doi:10.1242/jeb. 188581
- Barshis, D. J., J. H. Stillman, R. D. Gates, R. J. Toonen, L. W. Smith, and C. Birkeland. 2010. Protein expression and genetic structure of the coral *Porites lobata* in an environmentally extreme Samoan back reef: Does host genotype limit phenotypic plasticity? Mol. Ecol. **19**: 1705–1720. doi: 10.1111/j.1365-294X.2010.04574.x
- Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using lme4. J. Stat. Softw. **67**: 1–48. doi:10.18637/jss.v067.i01
- Bay, R. A., and S. R. Palumbi. 2015. Rapid acclimation ability mediated by transcriptome changes in reef-building corals. Genome Biol. Evol. 7: 1602–1612. doi:10.1093/gbe/evv085
- Bellantuono, A. J., O. Hoegh-Guldberg, and M. Rodriguez-Lanetty. 2012. Resistance to thermal stress in corals without changes in symbiont composition. Proc. R. Soc. B Biol. Sci. 279: 1100–1107. doi:10.1098/rspb.2011.1780
- Berkelmans, R., and B. L. Willis. 1999. Seasonal and local spatial patterns in the upper thermal limits of corals on the inshore Central Great Barrier Reef. Coral Reefs **18**: 219– 228. doi:10.1007/s003380050186
- Brown, B. E. 1997. Coral bleaching: Causes and consequences. Coral Reefs **16**: S129–S138.
- Camp, E. F., D. J. Smith, C. Evenhuis, I. Enochs, D. Manzello,S. Woodcock, and D. J. Suggett. 2016. Acclimatization tohigh-variance habitats does not enhance physiological

tolerance of two key Caribbean corals to future temperature and pH. Proc. R. Soc. B Biol. Sci. **283**: 20160442. doi:10. 1098/rspb.2016.0442

- Chan, N. C. S., and S. R. Connolly. 2013. Sensitivity of coral calcification to ocean acidification: A meta-analysis. Glob. Chang. Biol. **19**: 282–290. doi:10.1111/gcb.12011
- Coles, S. L., and P. L. Jokiel. 1977. Effects of temperature on photosynthesis and respiration in hermatypic corals. Mar. Biol. **43**: 209–216. doi:10.1007/bf00402313
- Cornwall, C. E., S. Comeau, T. M. DeCarlo, B. Moore, Q. D'Alexis, and M. T. McCulloch. 2018. Resistance of corals and coralline algae to ocean acidification: Physiological control of calcification under natural pH variability. Proc. R. Soc. B Biol. Sci. 285: 20181168. doi:10.1098/rspb.2018. 1168
- Dandan, S. S., J. L. Falter, R. J. Lowe, and M. T. McCulloch. 2015. Resilience of coral calcification to extreme temperature variations in the Kimberley region, northwest Australia. Coral Reefs 34: 1151–1163. doi:10.1007/s00338-015-1335-6
- Dixon, G. B., S. W. Davies, G. A. Aglyamova, E. Meyer, L. K. Bay, and M. V. Matz. 2015. Genomic determinants of coral heat tolerance across latitudes. Science (80-) **348**: 1460–1462. doi:10.1126/science.1261224
- Eakin, C. M., and others. 2010. Caribbean corals in crisis: Record thermal stress, bleaching, and mortality in 2005. PLoS One **5**: e13969.
- Fitt, W., B. Brown, M. Warner, and R. Dunne. 2001. Coral bleaching: Interpretation of thermal tolerance limits and thermal thresholds in tropical corals. Coral Reefs **20**: 51–65. doi:10.1007/s003380100146
- Frölicher, T. L., E. M. Fischer, and N. Gruber. 2018. Marine heatwaves under global warming. Nature 560: 360–364. doi:10.1038/s41586-018-0383-9
- Garcia, H. E., and L. I. Gordon. 1992. Oxygen solubility in seawater: Better fitting equations. Limnol. Oceanogr. **37**: 1307–1312. doi:10.4319/lo.1992.37.6.1307
- Grottoli, A. G., L. J. Rodrigues, and J. E. Palardy. 2006. Heterotrophic plasticity and resilience in bleached corals. Nature **440**: 1186–1189. doi:10.1038/nature04565
- Horvath, K. M., K. D. Castillo, P. Armstrong, I. T. Westfield, T. Courtney, and J. B. Ries. 2016. Next-century ocean acidification and warming both reduce calcification rate, but only acidification alters skeletal morphology of reef-building coral Siderastrea siderea. Sci. Rep. 6: 1–12. doi:10.1038/ srep29613
- Hughes, T. P., and others. 2018. Global warming transforms coral reef assemblages. Nature **556**: 492–496. doi:10.1038/ s41586-018-0041-2
- Iglesias-Prieto, R., V. H. Beltrán, T. C. LaJeunesse, H. Reyes-Bonilla, and P. E. Thomé. 2004. Different algal symbionts explain the vertical distribution of dominant reef corals in the eastern Pacific. Proc. R. Soc. B Biol. Sci. **271**: 1757– 1763. doi:10.1098/rspb.2004.2757

- Jokiel, P. L., J. E. Maragos, and L. Franzisket. 1978. Coral growth: Buoyant weight technique, p. 529–541. *In* D. R. Stoddart and R. E. Johannes [eds.], Coral reefs: Research methods. UNESCO.
- Jung, E. M. U., M. Stat, L. Thomas, A. Koziol, and V. Schoepf. 2021. Coral host physiology and symbiont dynamics associated with differential recovery from mass bleaching in an extreme, macro-tidal reef environment in northwest Australia. Coral Reefs 1–13: 893–905. doi:10.1007/s00338-021-02094-x
- Kapsenberg, L., and T. Cyronak. 2019. Ocean acidification refugia in variable environments. Glob. Chang. Biol. **25**: 3201–3214. doi:10.1111/gcb.14730
- Kenkel, C. D., S. P. Setta, and M. V. Matz. 2015. Heritable differences in fitness-related traits among populations of the mustard hill coral, *Porites astreoides*. Heredity (Edinb). **115**: 509–516. doi:10.1038/hdy.2015.52
- Klepac, C. N., and D. J. Barshis. 2020. Reduced thermal tolerance of massive coral species in a highly variable environment. Proc. R. Soc. B Biol. Sci. 287: 20201379. doi:10.1098/ rspb.2020.1379
- Kornder, N. A., B. M. Riegl, and J. Figueiredo. 2018. Thresholds and drivers of coral calcification responses to climate change. Glob. Chang. Biol. **24**: 5084–5095. doi:10.1111/ gcb.14431
- Kroeker, K. J., R. L. Kordas, R. Crim, I. E. Hendriks, L. Ramajo, G. S. Singh, C. M. Duarte, and J.-P. Gattuso. 2013. Impacts of ocean acidification on marine organisms: Quantifying sensitivities and interaction with warming. Glob. Chang. Biol. **19**: 1884–1896. doi:10.1111/gcb.12179
- LaJeunesse, T. C., J. E. Parkinson, P. W. Gabrielson, H. J. Jeong, J. D. Reimer, C. R. Voolstra, and S. R. Santos. 2018. Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. Curr. Biol. 28: 2570–2580. doi:10.1016/j.cub.2018.07.008
- Liu, G., and others. 2014. Reef-scale thermal stress monitoring of coral ecosystems: New 5-km global products from NOAA coral reef watch. Remote Sens. (Basel) **6**: 11579–11606. doi: 10.3390/rs61111579
- Mayfield, A. B., P. H. Chan, H. M. Putnam, C. S. Chen, and T. Y. Fan. 2012. The effects of a variable temperature regime on the physiology of the reef-building coral Seriatopora hystrix: Results from a laboratory-based reciprocal transplant. J. Exp. Biol. **215**: 4183–4195. doi:10.1242/jeb. 071688
- Mayfield, A. B., M.-N. Chen, P.-J. Meng, H.-J. Lin, C.-S. Chen, and P.-J. Liu. 2013. The physiological response of the reef coral Pocillopora damicornis to elevated temperature: Results from coral reef mesocosm experiments in Southern Taiwan. Mar. Environ. Res. **86**: 1–11. doi:10.1016/j. marenvres.2013.01.004
- Maynard, J. A., and others. 2008. ReefTemp: An interactive monitoring system for coral bleaching using high-resolution SST and improved stress predictors. Geophys. Res. Lett. **35**: L05603. doi:10.1029/2007gl032175

- Middlebrook, R., K. R. N. Anthony, O. Hoegh-Guldberg, and S. Dove. 2010. Heating rate and symbiont productivity are key factors determining thermal stress in the reef-building coral Acropora formosa. J. Exp. Biol. **213**: 1026–1034. doi: 10.1242/jeb.031633
- Middlebrook, R., K. R. N. Anthony, O. Hoegh-Guldberg, and S. Dove. 2012. Thermal priming affects symbiont photosynthesis but does not alter bleaching susceptibility in Acropora millepora. J. Exp. Mar. Bio. Ecol. **432–433**: 64–72. doi:10.1016/j.jembe.2012.07.005
- Muscatine, L., L. R. McCloskey, and R. E. Marian. 1981. Estimating the daily contribution of carbon from zooxanthellae to coral animal respiration1. Oceanography **26**: 601– 611. doi:10.4319/lo.1981.26.4.0601
- Le Nohaïc, M., C. L. Ross, C. E. Cornwall, S. Comeau, R. Lowe, M. T. McCulloch, and V. Schoepf. 2017. Marine heatwave causes unprecedented regional mass bleaching of thermally resistant corals in northwestern Australia. Sci. Rep. 7: 14999. doi:10.1038/s41598-017-14794-y
- Noonan, S. H. C., and K. E. Fabricius. 2015. Ocean acidification affects productivity but not the severity of thermal bleaching in some tropical corals. ICES J. Mar. Sci. **73**: 715– 726. doi:10.1093/icesjms/fsv127
- Oliver, E. C. J., and others. 2018. Longer and more frequent marine heatwaves over the past century. Nat. Commun. **9**: 1324. doi:10.1038/s41467-018-03732-9
- Oliver, T. A., and S. R. Palumbi. 2011. Do fluctuating temperature environments elevate coral thermal tolerance? Coral Reefs **30**: 429–440. doi:10.1007/s00338-011-0721-y
- Palumbi, S. R., D. J. Barshis, N. Traylor-Knowles, and R. A. Bay. 2014. Mechanisms of reef coral resistance to future climate change. Science (80-) 344: 895–898. doi:10.1126/science.1251336
- Putnam, H. M., and P. J. Edmunds. 2011. The physiological response of reef corals to diel fluctuations in seawater temperature. J. Exp. Mar. Bio. Ecol. **396**: 216–223. doi:10.1016/ j.jembe.2010.10.026
- R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Rivest, E. B., S. Comeau, and C. E. Cornwall. 2017. The role of natural variability in shaping the response of coral reef organisms to climate change. Curr. Clim. Change Rep. 3: 271–281. doi:10.1007/s40641-017-0082-x
- Safaie, A., and others. 2018. High frequency temperature variability reduces the risk of coral bleaching. Nat. Commun. 9: 1671. doi:10.1038/s41467-018-04074-2
- Schoepf, V., S. A. Carrion, S. M. Pfeifer, M. Naugle, L. Dugal, J. Bruyn, and M. McCulloch. 2019. Stress-resistant corals may not acclimatize to ocean warming but maintain heat

tolerance under cooler temperatures. Nat. Commun. **10**: 4031. doi:10.1038/s41467-019-12065-0

- Schoepf, V., C. E. Cornwall, S. M. Pfeifer, S. A. Carrion, C. Alessi, S. Comeau, and M. T. McCulloch. 2018. Impacts of coral bleaching on pH and oxygen gradients across the coral concentration boundary layer: A microsensor study. Coral Reefs 37: 1169–1180. doi:10.1007/s00338-018-1726-6
- Schoepf, V., M. U. Jung, M. T. McCulloch, N. White, M. Stat, and L. Thomas. 2020. Thermally variable, macrotidal reef habitats promote rapid recovery from mass coral bleaching. Front. Mar. Sci. 7: 245. doi:10.3389/FMARS.2020.00245
- Schoepf, V., M. Stat, J. L. Falter, and M. T. McCulloch. 2015. Limits to the thermal tolerance of corals adapted to a highly fluctuating, naturally extreme temperature environment. Sci. Rep. 5: 17639. doi:10.1038/srep17639
- Siebeck, U. E., N. J. Marshall, A. Klüter, and O. Hoegh-Guldberg. 2006. Monitoring coral bleaching using a colour reference card. Coral Reefs 25: 453–460. doi:10.1007/ s00338-006-0123-8
- Wall, C. B., T. Y. Fan, and P. J. Edmunds. 2013. Ocean acidification has no effect on thermal bleaching in the coral Seriatopora caliendrum. Coral Reefs 1–12: 119–130. doi:10. 1007/s00338-013-1085-2
- Warner, M. E., W. K. Fitt, and G. W. Schmidt. 1999. Damage to photosystem II in symbiotic dinoflagellates: A determinant of coral bleaching. Proc. Natl. Acad. Sci. USA 96: 8007–8012. doi:10.1073/pnas.96.14.8007
- van der Zande, R. M., M. Achlatis, D. Bender-Champ, A. Kubicek, S. Dove, and O. Hoegh-Guldberg. 2020. Paradise lost: End-of-century warming and acidification under business-as-usual emissions have severe consequences for symbiotic corals. Glob. Chang. Biol. 26: 2203–2219. doi:10. 1111/gcb.14998

### Acknowledgments

We thank the Bardi Jawi people who enabled this research through their advice and consent to access their traditional lands; James Brown, Gary Firman, and the staff at Cygnet Bay Pearl Farm for assistance in the field; and Prof. Malcolm McCulloch and the Australian Research Council Centre of Excellence for Coral Reef Studies for providing funding.

### **Conflict of Interest**

None declared.

Submitted 09 June 2021 Revised 27 September 2021 Accepted 05 December 2021

Associate editor: David Michael Baker

Coral heat tolerance under variable temperatures: Effects of different variability regimes and past environmental history versus current exposure

Running head: Coral tolerance to variable heat stress

Verena Schoepf<sup>1,2,3\*</sup>, Hermione Sanderson<sup>2,3,4</sup>, Ellis Larcombe<sup>2,3,4,5</sup>

<sup>1</sup>Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

<sup>2</sup>Oceans Graduate School and UWA Oceans Institute, The University of Western Australia, 35 Stirling Highway, Perth WA 6009, Australia

<sup>3</sup>ARC Centre of Excellence for Coral Reef Studies, The University of Western Australia, 35 Stirling Highway, Perth WA 6009, Australia

<sup>4</sup>Department of Biosciences, Swansea University, Swansea, SA2 8PP, United Kingdom

<sup>5</sup>Present address: Institute of Biomedical and Environmental Health, School of Health and Life Sciences, University of the West of Scotland, Paisley, United Kingdom

\*Corresponding author: Verena Schoepf@uva.nl)

Orcid ID Verena Schoepf: 0000-0002-9467-1088

Orcid ID Ellis Larcombe: 0000-0002-1609-6954

Keywords: coral heat tolerance, environmental history, environmental variability, temperature, light, physiology

# **Supplemental Information**

# **Additional Methods**

# Recovery of heat-stressed corals during the pre-conditioning phase

During the pre-conditioning phase, some colonies (C, D, E) were exposed to elevated temperatures (~32.4°C) from 17-30 May 2017 as part of the stress test in Schoepf et al. (2019), whereas others (A, B) served as ambient controls. Immediately after the stress test, temperature was returned to ambient seasonal temperatures, thus allowing the heat-stressed corals to recover for more than 8 months (1 June 2017 – 5 February 2018). Coral photochemical efficiency, calcification and visual health data show that they were fully recovered after 2, 4 and 8 weeks of recovery, respectively, when their values were no longer significantly different from the colonies that were not exposed to elevated temperatures (see below for details).

Coral photochemical efficiency (Fv/Fm), visual health and calcification rates were recorded every 2-4 weeks to monitor recovery following the methods described in the main text. t-tests were used to determine when values of recovering corals were no longer different from those of the ambient controls, i.e. by when they could be considered fully recovered with respect to these variables. Normality and homogeneity of variance was checked using the Shapiro-Wilk and F-test, respectively. When both assumptions were met, Student's t-test was used whereas the Welch's test was used when variances were not equal. When neither assumption was met, the non-parametric Wilcoxon test was used. All statistical analyses were run in R software.

Photochemical efficiency was already fully recovered 2 weeks after the heat stress test as Fv/Fm was no longer significantly different between control and recovering corals (t = 1.6904, df

= 5, p-value = 0.1517). Calcification rates measured 4 weeks after the heat stress test also showed no significant difference between control and recovering corals (Student's t = 0.37308, df = 5, p-value = 0.7244). Coral visual health was still significantly lower (-22%) in recovering compared to control corals 6 weeks after the heat stress test (Welch's t = 4.4907, df = 4, p-value = 0.0109) but that was no longer the case after 8 weeks (Wilcoxon's W = 7.5, p-value = 0.2059). Thus, with respect to these measurements, corals were fully recovered within 2-8 weeks after the heat stress test, while the pre-conditioning phase continued for another 6 months after this time period. Nevertheless, we acknowledge that other aspects of their physiology may not have been fully recovered, though we consider this unlikely given that (i) elevated temperatures during the stress test were only ~0.6°C above their bleaching threshold, (ii) heat stress only lasted for 2 weeks, (iii) recovery lasted for more than 8 months and (iv) corals were regularly fed throughout this period which has been shown to enhance recovery (e.g. Connolly et al. 2012).

### **Temperature variability experiment**

# Tidal light regime

From 2 Feb to 28 March 2018, light levels were 1  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup> at 06:00, 300 at 08:00, 560 at 11:00, 300 at 11:30 and 0 at 18:00, with gradual increases/decreases between set points. Light intensity in the morning and afternoon (but not maximum light levels at 11am) was then slightly increased from 29 March onwards for the rest of the temperature variability experiment and heat stress test (1  $\mu$ mol m<sup>-2</sup> s<sup>-1</sup> at 06:00, 400 at 08:00, 560 at 11:00, 400 at 11:30 and 0 at 18:00; Fig. 2b).

# Monitoring of experimental conditions

Due to logger malfunctioning, temperature was not recorded in one tank from 10 April – 10 May 2018 (constant heat-stress replicate tank #2). Therefore, during this time, we used data from the Apex temperature probe (10 min logging interval) for this tank to calculate treatment averages (Table 1).

From the acclimation phase onwards, pH<sub>T</sub>, salinity and temperature were measured at ~weekly time intervals in each tank around noon using a Schott Handylab pH 12 electrode (SI Analytics, Xylem Analytics, Germany; calibrated using TRIS buffer) and YSI 85 salinity meter (YSI, Yellow Springs, OH, USA), respectively. Water samples were also collected for total alkalinity (A<sub>T</sub>) analyses. These measurements revealed that pH (total scale) in the tanks was somewhat lower ( $\sim 7.8 - 7.9$ ) than ambient seawater pH but similar across treatments since all tanks received the same water (Table S3). Several independent studies have shown that bleaching sensitivity of 6 different coral species (including Acropora spp.) is not affected by seawater pH (Wall et al. 2013, 2018; Noonan and Fabricius 2015; Horvath et al. 2016; van der Zande et al. 2020); thus, the effect of the different temperature variability regimes on coral heat tolerance tested here should be independent of seawater pH. Water samples for alkalinity were refrigerated at ~2°C and analysed within 3 days or less. A<sub>T</sub> of water samples and certified reference materials provided by A.G. Dickson was analysed using potentiometric titration on a Mettler Toledo T50 titrator (Greifensee, Switzerland) and calculated using a modified Gran function, as described in Dickson et al. (2007). A<sub>T</sub>, pH<sub>T</sub>, temperature and salinity were then used to calculate seawater carbonate chemistry using the seacarb package in R and the constants recommended by Dickson et al. (2007).

# Statistical analyses

For LME models, significance of fixed effects were determined using Type-III sum of squares with Satterthwaite approximate of degrees of freedom from the *lmerTest* package (Kuznetsova et al. 2017). Significance of random effects was assessed using the ranova() function in *lmerTest* whereby each random-effect term is reduced or removed and likelihood ratio tests of model reductions are presented. For GLMM models, significance of fixed effects was determined using the Anova() function from the *car* package (Fox and Weisberg 2019), which provided Analysis of Deviance tables (Type III Wald chisquare tests). Since these are known to be anti-conservative, a stricter significance level of p≤0.01 was adopted for GLMM models. Pairwise contrasts were conducted using the *emmeans* package (Lenth 2021) when main effects and/or interaction terms were significant, with Tukey adjusted *p*-levels and the Kenward-Roger degrees-of-freedom method. Assumptions of normality and homoscedasticity for LME models were assessed using visual assessment of the residuals.

# References

- Connolly, S. R., M. A. Lopez-Yglesias, and K. R. N. Anthony. 2012. Food availability promotes rapid recovery from thermal stress in a scleractinian coral. Coral Reefs **31**: 951–960. doi:10.1007/s00338-012-0925-9
- Dandan, S. S., J. L. Falter, R. J. Lowe, and M. T. McCulloch. 2015. Resilience of coral calcification to extreme temperature variations in the Kimberley region, northwest Australia. Coral Reefs 34: 1151–1163. doi:10.1007/s00338-015-1335-6
- Dickson, A. G., C. L. Sabine, and J. R. Christian. 2007. Guide to Best Practices for Ocean CO2 Measurements. PICES Spec. Publ. 3: 191pp.
- Fox, J., and S. Weisberg. 2019. An R Companion to Applied Regression, Third. Sage.
- Horvath, K. M., K. D. Castillo, P. Armstrong, I. T. Westfield, T. Courtney, and J. B. Ries. 2016. Next-century ocean acidification and warming both reduce calcification rate, but only acidification alters skeletal morphology of reef-building coral Siderastrea siderea. Sci. Rep. 6: 1–12. doi:10.1038/srep29613
- Jung, E. M. U., M. Stat, L. Thomas, A. Koziol, and V. Schoepf. 2021. Coral host physiology and symbiont dynamics associated with differential recovery from mass bleaching in an extreme, macro-tidal reef environment in northwest Australia. Coral Reefs 1–13. doi:10.1007/s00338-021-02094-x
- Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. ImerTest Package: Tests in Linear Mixed Effects Models . J. Stat. Softw. 82: 1–26. doi:10.18637/jss.v082.i13
- Lenth, R. V. 2021. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.5.4.
- Le Nohaïc, M., C. L. Ross, C. E. Cornwall, S. Comeau, R. Lowe, M. T. McCulloch, and V.

Schoepf. 2017. Marine heatwave causes unprecedented regional mass bleaching of thermally resistant corals in northwestern Australia. Sci. Rep. 7: 14999. doi:10.1038/s41598-017-14794-y

- Noonan, S. H. C., and K. E. Fabricius. 2015. Ocean acidification affects productivity but not the severity of thermal bleaching in some tropical corals. ICES J. Mar. Sci. J. du Cons. doi:10.1093/icesjms/fsv127
- Schoepf, V., S. A. Carrion, S. M. Pfeifer, M. Naugle, L. Dugal, J. Bruyn, and M. McCulloch. 2019. Stress-resistant corals may not acclimatize to ocean warming but maintain heat tolerance under cooler temperatures. Nat. Commun. **10**: 4031. doi:10.1038/s41467-019-12065-0
- Schoepf, V., J. P. D'Olivo, C. Rigal, E. M. U. Jung, and M. T. McCulloch. 2021. Heat stress differentially impacts key calcification mechanisms in reef-building corals. Coral Reefs. doi:10.1007/s00338-020-02038-x
- Schoepf, V., E. M. U. Jung, M. McCulloch, N. E. White, M. Stat, and L. Thomas. 2020. Differential recovery from mass coral bleaching on naturally extreme reef environments in NW Australia. MarXiv. doi:10.31230/OSF.IO/S9XHA
- Schoepf, V., M. Stat, J. L. Falter, and M. T. McCulloch. 2015. Limits to the thermal tolerance of corals adapted to a highly fluctuating, naturally extreme temperature environment. Sci. Rep. 5: 17639. doi:10.1038/srep17639
- Wall, C. B., T. Y. Fan, and P. J. Edmunds. 2013. Ocean acidification has no effect on thermal bleaching in the coral Seriatopora caliendrum. Coral Reefs 1–12. doi:10.1007/s00338-013-1085-2

Wall, C. B., C. A. Ricci, G. E. Foulds, L. D. Mydlarz, R. D. Gates, and H. M. Putnam. 2018. The

effects of environmental history and thermal stress on coral physiology and immunity. Mar. Biol. **165**: 56. doi:10.1007/s00227-018-3317-z

van der Zande, R. M., M. Achlatis, D. Bender-Champ, A. Kubicek, S. Dove, and O. Hoegh-Guldberg. 2020. Paradise lost: End-of-century warming and acidification under business-asusual emissions have severe consequences for symbiotic corals. Glob. Chang. Biol. gcb.14998. doi:10.1111/gcb.14998

# **Supplementary Tables**

Table S1: Monthly seasonal sea surface temperature (SST) data for the Kimberley regionin Western Australia. The SST data served as target temperatures during the pre-conditioningphase. Data sourced from the U.S. National Oceanic and Atmospheric Administration (NOAA),Coral Reef Watch (version 2), 5-km virtual station "North Western Australia".

Month	SST (°C)
January	30.1
February	30.2
March	30.6
April	30.8
May	29.7
June	27.9
July	26.1
August	25.6
September	26.5
October	28.0
November	29.3
December	29.6
Average	28.7

Table S2: Daily temperature variability regimes and hourly temperature offset (°C) of the variable treatments relative to the constant control. Following the principle of degree heating days (see Methods), cumulative hourly heat stress exposure was calculated as "degree heating hours" (DHH) where all positive hourly temperature anomalies (i.e. those exceeding the control temperature of ~30°C, highlighted in bold) were summed up. Treatments were designed so that they had almost identical average daily temperatures as well as cumulative hourly heat stress exposure (DHH) for the variable and tidal treatments.

Time	constant	variable	tidal	offset - variable	e offset - tidal
05:00	30	28	30	-2	0
06:00	30	29	31.5	-1	1.5
07:00	30	29	31.5	-1	1.5
08:00	30	29	31.5	-1	1.5
09:00	30	30	34	0	4
10:00	30	30	34	0	4
11:00	30	30	30	0	0
12:00	30	31	30	1	0
13:00	30	31	30	1	0
14:00	30	31	30	1	0
15:00	30	32	30	2	0
16:00	30	32	30	2	0
17:00	30	32	30	2	0
18:00	30	31	30	1	-1
19:00	30	31	29	1	-1
20:00	30	31	29	1	-1
21:00	30	30	29	0	-1
22:00	30	30	29	0	0
23:00	30	30	30	0	0
00:00	30	29	30	-1	0
01:00	30	29	30	-1	0
02:00	30	29	30	-1	0
03:00	30	28	30	-2	0
04:00	30	28	30	-2	0
Average	30.00	30.00	30.35		
DHH				12.00	12.50

# **Table S3: Seawater carbonate chemistry based on discrete water samples.** Mean ± s.e.m.

The number of measurements (n) varied between experimental phases due to their different duration. Data from replicate tanks were averaged for each sampling day and experimental phase.  $A_T = \text{total alkalinity}, \Omega_{arag} = \text{aragonite saturation state}.$ 

		n	Temp.	Salinity	рНт	AT	pCO <sub>2</sub>	$\Omega_{ m arag}$
			(°C)	(ppt)		(µmol kg <sup>-1</sup> )	(µatm)	
(a) acclin	nation pha	ase						
constant		16	30.24	37.06	7.85	2363	689	2.97
			$\pm 0.11$	$\pm 0.05$	$\pm 0.01$	$\pm 2.6$	±12	$\pm 0.04$
variable		16	30.50	37.06	7.84	2363	712	2.93
			±0.16	$\pm 0.04$	$\pm 0.004$	$\pm 2.9$	$\pm 9$	$\pm 0.03$
tidal		16	30.33	37.04	7.84	2362	705	2.93
			$\pm 0.08$	$\pm 0.05$	$\pm 0.01$	$\pm 2.7$	$\pm 10$	$\pm 0.03$
(b) temp	erature ra	mp-up	phase					
constant	control	2	29.90	36.90	7.83	2356	726	2.81
			$\pm 0.30$	$\pm 0.00$	$\pm 0.01$	$\pm 0.2$	$\pm 29$	$\pm 0.05$
	heated	2	31.20	36.90	7.82	2358	746	2.89
			$\pm 0.20$	$\pm 0.00$	$\pm 0.02$	$\pm 0.3$	$\pm 50$	$\pm 0.12$
variable	control	2	30.20	36.95	7.84	2356	706	2.89
			$\pm 0.20$	$\pm 0.05$	$\pm 0.01$	±1.5	±23	$\pm 0.08$
	heated	2	30.90	36.90	7.81	2354	753	2.83
			$\pm 0.30$	$\pm 0.00$	$\pm 0.002$	$\pm 0.7$	$\pm 4$	$\pm 0.04$
tidal	control	2	30.45	36.90	7.81	2360	760	2.78
			$\pm 0.05$	$\pm 0.10$	$\pm 0.01$	$\pm 0.3$	±23	$\pm 0.06$
	heated	2	30.75	36.90	7.80	2357	777	2.76
			$\pm 0.05$	$\pm 0.00$	$\pm 0.01$	$\pm 1.0$	±17	$\pm 0.05$
(c) heat s	stress test	*						
constant	control	4*	30.58	36.63	7.92	2340	518	3.56
		4	$\pm 0.28$	$\pm 0.11$	$\pm 0.04$	$\pm 9.0$	$\pm 50$	$\pm 0.21$
	heated	4*	32.33	36.60	7.89	2340	550	3.67
			$\pm 0.35$	$\pm 0.08$	$\pm 0.03$	$\pm 3.8$	$\pm 55$	$\pm 0.22$
variable	control	$4^*$	30.18	36.53	7.91	2339	563	3.33
			$\pm 0.11$	$\pm 0.09$	$\pm 0.02$	$\pm 6.8$	±42	±0.12
	heated	$4^*$	32.83	36.65	7.89	2350	586	3.58
			$\pm 0.20$	$\pm 0.10$	$\pm 0.03$	$\pm 8.7$	$\pm 57$	±0.24
tidal	control	$4^*$	29.95	36.48	7.91	2341	554	3.37
			$\pm 0.09$	$\pm 0.09$	$\pm 0.04$	$\pm 6.5$	$\pm 63$	±0.23
	heated	$4^*$	32.50	36.55	7.89	2335	535	3.56
			$\pm 0.26$	$\pm 0.03$	$\pm 0.03$	n/a	n/a	n/a

\* temp, salinity, pH: n=4; all other parameters: CC n=3, CH n=2, VC n=3, VH n=3, TC n=3, TH n=1

Table S4: Results from (generalized) linear mixed models testing for the effects of variability regime and time on health score, Fv/Fm and Qm of *Acropora aspera* during the acclimation phase. Parent colony and fragment ID were included as random factors. var (fixed effects) = variability regime. Df = degrees of freedom. *Var*. (random effects) = variance. SD = Standard deviation. SS = sum of squares. MS = mean squares. For LME random effects, the % of the residual variance explained by the random effect is indicated in the posthoc column. Npar = number of model parameters. LRT = likelihood ratio test statistic. Significant p-values are highlighted in bold.

GLMM MODI	ELS			Df	Chisq	p(>Chisq)	Posthoc			
Health (GLMM, Gamma, Link = identity)										
var				2	0.68	0.7128				
time				1	1.76	0.1849				
var:time				2	0.25	0.8846				
Random effects		Var.	SD							
colony	intercept	0.07	0.27							
fragment ID	intercept	0.06	0.24							
residual	1	< 0.01	< 0.01							
LME										
MODELS	SS	MS	NumDF	DenDF	F	p(>F)	Posthoc			
Fv/Fm (LME)										
var	0.002	0.001	2	34.89	2.77	0.0764				
time	0.035	0.018	2	78.00	64.48	<0.0001	all sign.			
							different			
var:time	0.003	0.001	4	78.00	3.13	0.0193	see text			
random effect	npar	logLik	AIC	LRT	Df	p(>Chisq)				
none	12	262.34	-500.69							
colony	11	254.47	-486.95	15.74	1	0.0001	48%			
fragment ID	11	240.53	-459.05	43.63	1	<0.0001	33%			
Qm (LME)										
var	0.009	0.004	2	34.98	3.52	0.0406	constant >			
							variable			

time	0.002	0.001	2	78.00	0.75	0.4776	
var:time	0.007	0.002	4	78.00	1.38	0.2475	
random effect	npar	logLik	AIC	LRT	Df	p(>Chisq)	
none	12	187.70	-351.40				
colony	11	180.60	-339.20	14.20	1	0.0002	37%
fragment ID	11	178.69	-335.38	18.02	1	<0.0001	27%

Table S5: Results from (generalized) linear mixed models testing for the effects of variability regime, heat and time on health score, Fv/Fm, Qm and calcification rate of *Acropora aspera* during the temperature ramp-up and heat stress phase. Model structure varied slightly for each response variable (see Methods). var (fixed effects) = variability regime. Df = degrees of freedom. *Var*. (random effects) = variance. SD = Standard deviation. SS = sum of squares. MS = mean squares. For LME random effects, the % of the residual variance explained by the random effect is indicated in the posthoc column. Npar = number of model parameters. LRT = likelihood ratio test statistic. Significant p-values are highlighted in bold.

# GLMM MODELS Df Chisq p(>Chisq) Posthoc

Health - heat stress phase (GLMM, Gamma, link=identity)									
heat				1	0.11	0.7404			
var				2	11.06	0.0040	none sign.		
heat:var				2	12.60	0.0018	see text		
Random									
effects		Var.	SD						
colony	intercept	0.11	0.34						
time	intercept	0.03	0.17						
residual		0.02	0.15						

# Health after 9 days of heat stress (day 108) (GLMM, Gamma, link=identity)

heat				1	2.02	0.1550	
var				2	61.51	<0.0001	constant = variable > tidal
heat:var				2	42.81	<0.0001	see text
Random							
effects		Var.	SD				
colony	intercept	0.21	0.45				
residual		0.01	0.11				

# Health 3 days after end of heat stress test (day 113) (GLMM, Gamma, link=identity)

heat	1	0.60	0.4386	
var	2	15.28	0.0005	constant ≠ tidal
heat:var	2	10.77	0.0046	see text

Random			
effects		Var.	SD
colony	intercept	0.21	0.46
residual		0.03	0.18

# Fv/Fm - heat stress phase (GLMM, Gamma, link=identity)

var				2	68.76	<0.0001	constant = tidal > variable
heat				1	0.87	0.3517	
var:heat				2	45.53	<0.0001	see text
Random							
effects		Var.	SD				
colony	intercept	< 0.01	0.01				
time	intercept	< 0.01	0.01				
residual		< 0.01	0.05				

# Fv/Fm after 10 days of heat stress (day 109) (GLMM, Gamma, link=identity)

var	·		, .	2	64.21	<0.0001	constant > variable = tidal
heat				1	0.001	0.9753	
var:heat				2	46.76	<0.0001	see text
Random							
effects		Var.	SD				
colony	intercept	< 0.01	0.02				
residual		< 0.01	0.05				

# Fv/Fm after 11 days of heat stress (day 110) (GLMM, Gamma, link=identity)

var				2	22.90	<0.0001	constant > variable = tidal
heat				1	0.026	0.8720	
var:heat				2	12.58	0.0019	see text
Random							
effects		Var.	SD				
colony	intercept	< 0.01	0.03				
residual		0.01	0.11				
LME							

MODELS	SS	MS	NumDF	DenDF	F	p(>F)	Posthoc	
Qm - heat stress	phase (I	LME)						
var	0.08	0.04	2	29.41	8.19	0.0015	variable ≠ tidal	
heat	0.01	0.01	1	29.73	1.36	0.2530		
time	0.15	0.02	7	157.89	4.27	0.0002	see text	

	0.05	0.02	2	20.50	4 00	0.0126	
var:neat	0.05	0.03	2	29.50	4.99	0.0130	see text
var:time	0.14	0.01	14	157.53	1.98	0.0223	see text
heat:time	0.21	0.03	7	157.90	5.91	<0.0001	see text
var:heat:time	0.10	0.01	14	157.60	1.46	0.1323	
random							
effects	npar	logLik	AIC	LRT	Df	p(>Chisq)	
none	51	160.69	-219.38				
colony	50	160.61	-221.22	0.16	1	0.6936	3%
fragment ID	50	127.08	-154.17	67.21	1	<0.0001	55%
Calcification -	entire exp	eriment (	LME)				
var	0.45	0.23	2	31.87	7.27	0.0025	constant = variable > tidal
heat	0.35	0.35	1	31.87	11.24	0.0021	control > heated
time	0.55	0.55	1	36.00	17.50	0.0002	acclim > stress
var:heat	0.17	0.08	2	31.87	2.65	0.0865	
var:time	0.32	0.16	2	36.00	5.18	0.0105	see text
heat:time	0.78	0.78	1	36.00	25.04	<0.0001	see text
var:heat:time	0.19	0.10	2	36.00	3.08	0.0584	
random							
effects	npar	logLik	AIC	LRT	Df	p(>Chisq)	
none	15	-7.49	44.97				
colony	14	-13.89	55.77	12.80	1	0.0003	42%
fragment ID	14	-10.04	48.08	5.10	1	0.0239	22%

Table S6: Results from linear mixed effect models testing for the effects of variability regime and heat on photosynthesis (P) and respiration (R) rate as well as P/R ratios of *Acropora aspera*. Parent colony was included as random factor. Var. = variability regime. SS = sum of squares. MS = mean squares. Df = degrees of freedom. Significant p-values are highlighted in bold. For random effects, the % of the residual variance explained by the random effect is indicated in the posthoc column. Npar = number of model parameters. LRT = likelihood ratio test statistic.

	SS	MS	NumDF	DenDF	F	p(>F)	Posthoc
Photosynthesis							
var	0.33	0.16	2	29.15	2.28	0.1205	
heat	2.95	2.95	1	29.18	41.01	<0.0001	control >
							heated
var:heat	0.37	0.18	2	29.15	2.56	0.0946	
random	npar	logLik	AIC	LRT	Df	p(>Chisq)	
effect	0				0		
none	8	-11.92	39.847	0.00	0	0.0000	
colony	7	-14.56	43.129	5.28	1	0.0216	32%
Respiration							
var	0.01	0.01	2	29.02	1.19	0.3200	
heat	0.00	0.00	1	29.06	0.11	0.7379	
var:heat	0.04	0.02	2	29.02	3.49	0.0438	None sign.
random	npar	logLik	AIC	LRT	Df	p(>Chisq)	
effect							
none	8	30.93	-45.852	0.00	0	0.0000	
colony	7	29.45	-44.908	2.94	1	0.0862	25%
P/R ratio							
var	3.83	1.92	2	29.36	4.41	0.0212	constant >
							tidal
heat	51.82	51.82	1	29.40	119.23	<0.0001	control >
	1.00				1.10	0.0010	heated
var:heat	1.03	0.51	2	29.36	1.18	0.3213	
random	npar	logLik	AIC	LRT	Df	p(>Chisq)	
effect	0	41.07	00.10	0.00	0	0.0000	
none	8	-41.06	98.12	0.00	0	0.0000	<b>25</b> 0 (
colony	7	-42.99	99.97	3.86	1	0.0496	25%

Table S7: Overview of research published to date to investigate the effects of temperature variability on coral physiology and heat tolerance in the macrotidal Kimberley region in NW Australia. The effects of temperature variability were investigated by comparing (1) corals from intertidal and subtidal habitats (high vs moderate daily temperature variability, respectively), or (2) different temperature variability regimes in the aquarium-based experiments. Studies highlighted in green indicate a positive effect of highly variable temperatures on coral heat tolerance, blue indicates no effect and red indicates a negative effect. Studies without highlighting demonstrated more complex effects.

Type of study	Main finding
(a) Field observations	
Dandan et al. 2015: Seasonal in	Massive Dipsastraea favus and Trachyphyllia geoffroyi
situ calcification rates for three	calcified faster in the intertidal/intermediate habitat,
coral species from three habitats	whereas branching Acropora aspera calcified faster in
(intertidal <sup>1</sup> , intermediate, subtidal)	the subtidal
Le Nohaïc et al. 2017: Coral	Both intertidal and subtidal coral communities suffered
community composition and health	from extensive bleaching, but the subtidal coral
status before and during the 2016	community had a greater percentage of severely
mass bleaching event (intertidal,	bleached corals
subtidal)	
Schoepf et al. 2020: Coral	The intertidal coral community was fully recovered six
community composition and health	months after the 2016 mass bleaching event, whereas
status six months after the 2016	the subtidal community suffered extensive mortality

mass bleaching event (intertidal,	
subtidal)	
Schoepf et al. 2021: Coral	The biomineralization response was generally highly
calcification mechanisms and	sensitive to heat stress but did not differ between
skeletal trace element composition	thermally distinct reef habitats (intertidal vs subtidal)
of A. aspera during a heatwave and	
"normal" year (intertidal, subtidal)	
Jung et al. 2021: Energy reserves,	Subtidal corals were more severely bleached than
symbiont community composition,	intertidal corals and suffered greater mortality.
cell density and chlorophyll a	Cladocopium dominated all corals, but symbiont
concentration of A. aspera during	community composition differed significantly between
and six months after the 2016 mass	environments and between bleached and healthy
bleaching event (intertidal,	subtidal corals. Bleaching resilience was decoupled
subtidal)	from energy reserve levels/catabolization.
(b) Experimental (aquarium-based	) studies
Schoepf et al. 2015: Aquarium-	Intertidal corals of both species (A. aspera, Dipsastraea
based heat stress test comparing the	sp.) had higher heat tolerance and survival than subtidal
heat tolerance of two coral species	conspecifics
from two habitats (intertidal,	
subtidal)	
Schoepf et al. 2019: Long-term	Exposure to daily, symmetric temperature variability of
exposure of intertidal A. aspera to	4°C over 9 months lowered coral heat tolerance
warmer and cooler temperatures	compared to corals maintained at constant daily

under both constant and variable	temperatures. Long-term exposure to 3-6°C cooler
temperatures, followed by a heat	temperature did not affect heat tolerance. In contrast,
stress test (intertidal only)	corals were not able to acclimatize to +1°C warming
	over 6 months.
This study: Effects of short- and	Pre-conditioning to constant vs variable temperatures
long-term environmental history	for 1.5 years did not significantly impact coral
and two different variability	physiology and heat tolerance. In contrast,
regimes on heat tolerance of	environmental history experienced in the month prior to
intertidal A. aspera (intertidal only)	the heat stress test significantly influenced physiological
	responses, with corals exposed to both types of
	variability having lower heat tolerance.

<sup>1</sup>referred to as 'isolated' in Dandan et al. 2015

# **Supplementary Figures**



**Figure S1: Daily temperature variability during the pre-conditioning phase, i.e. prior to the temperature variability experiment.** Schematic of the temperature regimes in treatments with constant daily temperature versus 4°C daily variability from 1 August 2016 until 5 Feb 2018.